

# Top Flavor Changing Interactions with Neutral Higgs Bosons

Chung Kao  
University of Oklahoma



<sup>†</sup>Presented at the Brookhaven Forum 2015 (BF2015)  
at Brookhaven National Laboratory, Upton, NY, October 7-9, 2015.

# When the Higgs meets the Top


Altunkaynak, Hou, Kao, Kohda, and McCoy (2015);  
Chen, Hou, Kao, and Kohda, Phys. Lett. B 725 (2013) 378;  
Kao, Cheng, Hou, and Sayre, Phys. Lett. B 716 (2012) 225.

- Introduction and Motivation
- General Two Higgs Doublet Models
- When the Higgs Meets the Top
- The Discovery Potential at the LHC
- Conclusions

# Heavyweight Champion before July 4, 2012



# The New Runner-up



# Theoretical Values for FCNC Top Decays

ATLAS-PHYS-PUB-2013-012

Process	SM	QS	2HDM-III	FC-2HDM	MSSM
$t \rightarrow u\gamma$	$3.7 \cdot 10^{-16}$	$7.5 \cdot 10^{-9}$	—	—	$2 \cdot 10^{-6}$
$t \rightarrow uZ$	$8 \cdot 10^{-17}$	$1.1 \cdot 10^{-4}$	—	—	$2 \cdot 10^{-6}$
$t \rightarrow uH$	$2 \cdot 10^{-17}$	$4.1 \cdot 10^{-5}$	$5.5 \cdot 10^{-6}$	—	$10^{-5}$
$t \rightarrow c\gamma$	$4.6 \cdot 10^{-14}$	$7.5 \cdot 10^{-9}$	$\sim 10^{-6}$	$\sim 10^{-9}$	$2 \cdot 10^{-6}$
$t \rightarrow cZ$	$1 \cdot 10^{-14}$	$1.1 \cdot 10^{-4}$	$\sim 10^{-7}$	$\sim 10^{-10}$	$2 \cdot 10^{-6}$
$t \rightarrow cH$	$3 \cdot 10^{-15}$	$4.1 \cdot 10^{-5}$	$1.5 \cdot 10^{-3}$	$\sim 10^{-5}$	$10^{-5}$

# Introduction and Motivation

Das and Kao (1996)

- A special two Higgs doublet model explains why top quark is the most massive elementary particle by suggesting that it is the only fermion that couples to a Higgs doublet ( $\phi_2$ ) with a much larger VEV ( $v_2 \gg v_1$ ).
- This model leads to flavor changing neutral Higgs (FCNH) interactions and CP violation.
- Most LHC data are consistent with the Standard Model. FCNH interactions might lead to new physics beyond SM.

# A Special Higgs Model for the Top Quark

## 1 Introduction

In the Standard Model (SM) of electroweak interactions:

1. There is one Higgs doublet to generate mass for gauge bosons as well as for fermions. A neutral Higgs scalar ( $H^0$ ) remains after spontaneous symmetry breaking.
2. The top quark has a large mass because its Yukawa coupling with the  $H^0$  is large.<sup>†</sup>

In a special two Higgs doublet model, the top quark is much heavier than the other quarks and the leptons, because it is the only elementary fermion getting a mass from a much larger vacuum expectation value (VEV) of a second Higgs doublet.

This model has a few interesting features:

1. The ratio of the Higgs VEVs,  $\tan \beta \equiv |v_2|/|v_1|$ , is chosen to be large.
2. The Yukawa couplings of the lighter fermions are highly enhanced.
3. There are flavor changing neutral Higgs interactions.

---

<sup>†</sup>The mass of a fermion is equal to its Yukawa coupling with the  $H^0$  times the vacuum expectation value of the Higgs field,  $m = \lambda(v/\sqrt{2})$ .

### 3 Special Yukawa Interactions

We choose the Lagrangian density of Yukawa interactions to be of the following form

$$\begin{aligned}\mathcal{L}_Y = & - \sum_{m,n=1}^3 \bar{L}_L^m \phi_1 E_{mn} l_R^n - \sum_{m,n=1}^3 \bar{Q}_L^m \phi_1 F_{mn} d_R^n \\ & - \sum_{\alpha=1}^2 \sum_{m=1}^3 \bar{Q}_L^m \tilde{\phi}_1 G_{m\alpha} u_R^\alpha - \sum_{m=1}^3 \bar{Q}_L^m \tilde{\phi}_2 G_{m3} u_R^3 + \text{H.c.},\end{aligned}$$

where

$$\phi_\alpha = \begin{pmatrix} \phi_\alpha^+ \\ \frac{v_\alpha + \phi_\alpha^0}{\sqrt{2}} \end{pmatrix}, \quad \tilde{\phi}_\alpha = \begin{pmatrix} \frac{v_\alpha^* + \phi_\alpha^{0*}}{\sqrt{2}} \\ -\phi_\alpha^- \end{pmatrix}, \quad \phi_\alpha^- = \phi_\alpha^{+*}, \quad \alpha = 1, 2, \quad \text{and} \quad (5)$$

$$L_L^m = \begin{pmatrix} \nu_l \\ l \end{pmatrix}_L^m, \quad Q_L^m = \begin{pmatrix} u \\ d \end{pmatrix}_L^m, \quad m = 1, 2, 3, \quad (6)$$

$l^m$ ,  $d^m$ , and  $u^m$  are the gauge eigenstates.

This Lagrangian respects a discrete symmetry,

$$\begin{aligned}\phi_1 & \rightarrow -\phi_1, \quad \phi_2 \rightarrow +\phi_2, \\ l_R^m & \rightarrow -l_R^m, \quad d_R^m \rightarrow -d_R^m, \quad u_R^\alpha \rightarrow -u_R^\alpha, \\ L_L^m & \rightarrow +L_L^m, \quad Q_L^m \rightarrow +Q_L^m, \quad u_R^3 \rightarrow +u_R^3.\end{aligned} \quad (7)$$



# A General Two Higgs Doublet Model

Mahmoudi and Stal (2009)

- Let us express the general Yukawa interaction Lagrangian for neutral Higgs bosons as

$$\begin{aligned}\sqrt{2}\mathcal{L}_I^N = & \bar{U} [-\kappa^U s_{\beta-\alpha} - \rho^U c_{\beta-\alpha}] U h^0 + \bar{D} [-\kappa^D s_{\beta-\alpha} - \rho^D c_{\beta-\alpha}] D h^0 \\ & + \bar{U} [-\kappa^U c_{\beta-\alpha} + \rho^U s_{\beta-\alpha}] U H^0 + \bar{D} [-\kappa^D c_{\beta-\alpha} + \rho^D s_{\beta-\alpha}] D H^0 \\ & + \bar{U} [+i\gamma_5 \rho^U] U A^0 + \bar{D} [-i\gamma_5 \rho^D] D A^0\end{aligned}$$

where  $\kappa^f = \frac{\sqrt{2}m_f}{v}$ ,  $\tan \beta \equiv v_2/v_1$ , and  $v = \sqrt{v_1^2 + v_2^2}$ .

- There are 4 flavor conserving models with  $Z_2$  symmetries, such that  $\rho$ 's are related to  $\kappa$ 's in the following form [Barger, Hewett and Phillips, PRD **41** (1990) 3421.]:

	Type			
	I	II	III	IV
$\rho^D$	$\kappa^D \cot \beta$	$-\kappa^D \tan \beta$	$-\kappa^D \tan \beta$	$\kappa^D \cot \beta$
$\rho^U$	$\kappa^U \cot \beta$	$\kappa^U \cot \beta$	$\kappa^U \cot \beta$	$\kappa^U \cot \beta$
$\rho^E$	$\kappa^E \cot \beta$	$-\kappa^E \tan \beta$	$\kappa^E \cot \beta$	$-\kappa^E \tan \beta$

- In a general model without  $Z_2$  symmetries,  $\rho$  matrices are free.

# The Decoupling Limit of 2HDM

Gunion and Haber (2003)

- In the decoupling limit of 2HDM, we expect
  - ▶  $M_h = O(v)$
  - ▶  $M_H, M_A, M_{H^\pm} = M_S + O(v^2/M_S)$
  - ▶  $|\cos(\beta-\alpha)| = O(v^2/M_S^2)$
  - ▶ If  $\cos(\beta-\alpha) = 0$ ,  $h^0$  becomes the SM Higgs boson.
- Recently, there has been interests in the 2HDM parameter space where the alignment is obtained without decoupling and without fine tuning where  $H^0$  and  $A^0$  can be light and  $h^0$  is like SM Higgs.  
Craig, Galloway, Thomas (2013); Carena et al. (2014)

# Constraints on FCNH Couplings




- ATLAS and CMS data have placed tight constraints on  $\lambda_{tc}$  and  $\lambda_{ct}$  with  $t \rightarrow ch^0 \rightarrow c\gamma\gamma$ :
  - ▶ the top decay should have  $B(t \rightarrow ch^0) < 0.56\%$ ,
  - ▶ or  $\sqrt{\lambda_{tc}^2 + \lambda_{ct}^2} < 0.14$ , with  $\lambda_{ct} = \rho_{ct} \cos(\beta - \alpha)$ .
- If we choose  $\rho$ -matrix to be Hermitian, then  $b \rightarrow s\gamma$  and  $B - \bar{B}$  mixing imply  $|\rho_{ct}| < 0.1$ .
- If the  $\rho$ -matrix is not Hermitian, then we must have  $|\rho_{ct}| < 0.1$ , while  $|\rho_{tc}|$  can be close to 1.

# When the Higgs Meets the Top

- The Higgs boson is the mass giver, while the top quark is the most massive particle. Their interactions might give us guidance to search for new physics beyond the Standard Model.
- The LHC has become a top factory.
- We might be able to observe  $t \rightarrow ch^0$  if  $\lambda_{ct} = \rho_{ct} \cos(\beta - \alpha)$  can lead to observable signal.
- Or we might discover  $H^0, A^0 \rightarrow t\bar{c} + \bar{t}c$  in the decoupling limit with  $\lambda_{tc} = \rho_{tc} \sin(\beta - \alpha)$ .

# FCNH Top Decays at the LHC


Hou (1991); Hall and Weinberg (1993); Aguilar-Saavedra and Branco (2000);  
Kao, Cheng, Hou, and Sayre (2012); Chen, Hou, Kao, and Kohda (2013);  
Atwood, Gupta, Soni (2013).



$$g_{htc}(\text{CS}) = \frac{\sqrt{m_t m_c}}{v} \sim 0.06$$


$$\lambda_{htc}(\text{HW}) = \epsilon_{Q3} \epsilon_{U2} \sim 0.2$$

# Discovery Potential with 8 TeV



# Discovery Contours

$L = 20 \text{ fb}^{-1}$  at 8 TeV;  $30 \text{ fb}^{-1}$  at 14 TeV




# Constraint from the Golden Mode for Higgs Discovery

- The CMS preliminary result with full 7 and 8 TeV data shows 13, 8, and 4 events with 0, 1, and 2 jets, respectively, after selecting events with  $121.5 \text{ GeV} < M_{4l} < 130.5 \text{ GeV}$ .
- The resulting 95% confidence level limit on the relative signal strength between  $t$  to  $ch^0$  and inclusive Higgs production is around 31%,
- That can be converted to a limit of 6.5 pb on the effective cross section of  $t$  to  $ch^0$  at 8 TeV, or a branching ratio limit around 1.5%.



# The Golden Mode for Higgs Discovery




# Summary for FCNH top Decay


- It is of great interest to search for the link between the top quark ( $t$ ) and the Higgs bosons ( $H^0$ ,  $h^0$ ,  $A^0$ ).
- A discovery of  $t \rightarrow ch^0$  would suggest the existence of an extended Higgs sector beyond the usual 2HDM-II and MSSM.
- Experimental studies of  $h^0$  to  $bb$ ,  $WW^*$ ,  $ZZ^*$ ,  $\tau^+\tau^-$  and  $\gamma\gamma$  modes will provide important information for FCNH interactions.

# The FCNH Signal of a Heavy Higgs boson at the LHC



Let us consider a flavor changing neutral Higgs boson ( $\phi^0$ ) with  $M_\phi > M_h$ . It can be a CP-even scalar ( $H^0$ ) or a CP-odd pseudoscalar ( $A^0$ ) produced at the LHC followed by the Higgs decay into a top quark and a charm quark:




# Heavy Higgs Decay Branching Fractions



# Constraints from B Physics




# Signal versus Physics Background at the LHC with 8 TeV



# Physics Background

- $pp \rightarrow Wjj + X, j = u, d, s, c, \text{ or } g$
- $pp \rightarrow Wbb + X$
- Single top:



# Realistic Acceptance Cuts

We require that in every event there must be

- exactly 2 jets with  $p_T > 20$  GeV,  $|\eta| < 2.5$  and exactly one of the jets must be tagged as a b-jet;
- an isolated lepton with  $p_T > 20$  GeV,  $|\eta| < 2.5$ ;
- the missing transverse energy must be greater than 20 GeV;
- the angular separation between jets and the lepton must be  $\Delta R(b,j,l) > 0.4$ .




# Mass Reconstruction

We require that the reconstructed invariant masses should center around  $m_t$ ,  $m_W$ , and  $M_\phi$ .

- Assuming an on-shell  $W$ , we evaluate  $k_z$  of the neutrino with lepton momentum ( $p$ ) and missing transverse energy. Usually, there are two possible values for  $k_z$ . We select whichever leads to a better reconstruction of the top-quark mass:  $\text{Min}[m_t^2 - (k+p+p_b)^2]$ , and define the reconstructed top mass as  $M_t^R = M_{b\ell\nu}$  such that  $|M_{b\ell\nu} - m_t| < 0.15m_t$  or  $0.20m_t$ .
- The invariant mass of the top and the charm should have a peak near  $M_\phi$ :  $|M_{b\ell\nu j} - m_\phi| < 0.15M_\phi$  or  $0.20M_\phi$ .



# Discovery Contour at the LHC

## $\cos(\beta-\alpha) = 0.2$



# Discovery Contour at the LHC

## $\cos(\beta-\alpha) = 0.1$



# Conclusions

- It is of great interest to search for the link between the heaviest particle (top) and the mass giver (Higgs).
- It is a win-win strategy to search for the FCNH top decay  $t \rightarrow ch^0$  and the heavy Higgs decay  $H^0, A^0 \rightarrow t\bar{c} + t\bar{c}$ . In the decoupling limit, the production ( $gg \rightarrow H^0$ ) and the FCNH decay  $H^0 \rightarrow tc$  can be sustained by  $\sin(\beta-\alpha) \sim 1$ .
- The FCNH decay of heavy Higgs bosons will be observable for  $\rho_{tc} > 0.1$  and  $\cos(\beta-\alpha) \sim 0.1$  up to  $M_H = 800$  GeV with  $3000 \text{ fb}^{-1}$  of data.
- We might find out if nature chooses the same mechanism for electroweak symmetry breaking and tree-level FCNC.